

## Pneumococcal Conjugate Vaccine (PCV7) Requirements for Child-Care Attendance

Age of Child (Months)	Vaccination History	Required Doses
<b>2 – 6*</b>	0 doses	3 doses, 2 mos apart
	1 dose	2 doses, 2 mos apart
	2 doses	1 dose, 2 mos after the most recent dose
<b>7 – 11*</b>	0 doses	2 doses, 2 mos apart
	1 or 2 doses	1 dose, 2 mos after the most recent dose
<b>12 – 23</b>	0 doses	2 doses, $\geq 2$ mos apart
	1 dose administered before age 12 mos	2 doses, $\geq 2$ mos apart
	1 dose administered on or after age 12 mos	1 dose, $\geq 2$ mos after the most recent dose
	2 or 3 doses administered before age 12 mos	1 dose, $\geq 2$ mos after the most recent dose
<b>24 – 59</b>		
Healthy children	Any incomplete schedule <sup>†</sup>	1 dose, $>2$ mos after most recent dose
Children at high risk <sup>‡</sup>	Any incomplete schedule <sup>†</sup>	2 doses separated by 2 mos

\*Children in these age groups will need a booster dose on or after 12 months of age.

<sup>†</sup> Any of the scenarios for children aged 2-6 (with a booster on or after 12 months of age), 7-11 (with a booster on or after 12 months of age), and 12-23 months constitute a complete series.

<sup>‡</sup> Children with certain chronic conditions or immuno-suppression conditions will also require a dose of pneumococcal polysaccharide vaccine (PPV23) in addition to PCV7 two months after the last PCV7.